

10/573329

IA99 Rec'd PCT/PTO 24 MAR 2006

SPANISH PATENT AND TRADEMARK OFFICE

OFFICIAL CERTIFICATE

I hereby certify that the annexed documents are an exact copy of the PATENT OF INVENTION application number 200302643, that was filed before this Office on November 13, 2003.

Madrid, November 16, 2004

The Director of the Patents Department
and Technological Information

(signature)
CARLOS GARCIA NEGRETE

[Seal of the
Spanish Patents and
Trademarks Office]

SPANISH PATENTS AND TRADEMARKS OFFICE

[Stamp bearing: Files-Association of Industrial Property Agents]

APPLICATION NUMBER
P200302643

DATE AND HOUR OF FILING AT THE O.E.P.M.
13 November 2003

INSTANCE OF APPLICATION FOR:

☒ PATENT OF INVENTION

☐ UTILITY MODEL

(2) <input type="checkbox"/> ADDITIONAL APPLICATION <input type="checkbox"/> DIVISIONAL APPLICATION <input type="checkbox"/> CHANGE OF MODALITY <input type="checkbox"/> TRANSFORMATION <input type="checkbox"/> EUROPEAN APPLICATION	(3) PRINCIPAL OR ORIGINAL FILE MODALITY APPLICATION NUMBER FILING DATE / / MODALITY APPLICATION NUMBER FILING DATE / /	DATE AND HOUR OF FILING IN A PLACE OTHER THAN THE O.E.P.M. (4) PLACE OF FILING CODE MADRID 28
---	--	--

(5) APPLICANT (S) SURNAME OR JURIDICAL DENOMINATION	NAME	NATIONALITY	COUNTRY CODE	D.N.I.
QUÍMICA SINTETICA, S.A.		SPANISH	ES	A28008167

(5) PARTICULARS OF THE FIRST APPLICANT

ADDRESS: C. de l'Acer, 24

TOWN: BARCELONA

PROVINCE: BARCELONA

COUNTRY OF RESIDENCE: SPAIN

NATIONALITY: Spanish

PHONE | |

POST CODE 08004

COUNTRY CODE |ES|

NATION CODE |ES|

(7) INVENTOR (S)	(8) <input type="checkbox"/> THE APPLICANT IS THE INVENTOR <input checked="" type="checkbox"/> THE APPLICANT IS NOT THE INVENTOR OR SOLE INVENTOR	(9) WAY OF OBTENTION OF RIGHTS <input checked="" type="checkbox"/> EMPLOYER (S) <input type="checkbox"/> CONTRACT <input type="checkbox"/> SUCCESSION
------------------	--	--

SURNAMES	NAME	NATIONALITY	CODE
COSME GOMEZ	ANTONIO	SPANISH	ES
VILLASANTE PRIETO	JAVIER	SPANISH	ES
PALOMO NICOLAU	FRANCISCO	SPANISH	ES

(9) TITLE OF THE INVENTION	
----------------------------	--

CRYSTALLINE FORM OF GATIFLOXACIN

(11) MICROBIOLOGICAL DIPOSIT	<input type="checkbox"/> YES <input type="checkbox"/> NO
------------------------------	--

(12) OFFICIAL EXHIBITIONS: PLACE PLACE	DATE
--	------

(13) PRIORITY DECLARATIONS	
----------------------------	--

COUNTRY OF ORIGIN	CODE	NUMBER	DATE

(14) THE APPLICANTS REQUEST THE EXEMPTION OF THE PAYMENT OF TAXES PROVIDED IN ART. 162 L.P.	<input type="checkbox"/> YES <input type="checkbox"/> NO
---	--

(15) REPRESENTATIVE	SURNAMES: PONTI SALES	NAME CODE Adelaida 388/3
---------------------	-----------------------	------------------------------

ADDRESS C. Consell de Cent, 322	TOWN Barcelona	PROVINCE POSTAL CODE Barcelona 08007
------------------------------------	-------------------	---

(16) LIST OF ANNEXED DOCUMENTS	<input checked="" type="checkbox"/> SPECIFICATION N° OF PAGES 11 <input checked="" type="checkbox"/> CLAIMS N° OF PAGES 5 <input checked="" type="checkbox"/> DRAWINGS N° OF PAGES 3 <input type="checkbox"/> SEQUENCE LISTING 0 <input checked="" type="checkbox"/> ABSTRACT <input type="checkbox"/> PRIORITY DOCUMENT <input type="checkbox"/> TRANSLATION OF PRIORITY OCUMENT	<input checked="" type="checkbox"/> REPRESENTATION FORM <input type="checkbox"/> PROOFS <input checked="" type="checkbox"/> RECEIPT OF PAYMENT OF TAXES <input type="checkbox"/> SHEET OF COMPLEMENTARY. INFORMATION <input checked="" type="checkbox"/> OTHER Magnetic Support and Inventor's declaration.	SIGNATURE OF THE APPLICANT OR REPRESENTATIVE Adelaida Ponti Sales Associate no. 320 SIGNATURE OF THE OFFICIAL
--------------------------------	---	---	--

NOTIFICATION OF PAYMENT OF THE GRANT FEES

MINISTRY OF
 SCIENCE AND
 TECHNOLOGY

Spanish patents and trademarks Office

APPLICATION NUMBER

200302643

FILING DATE :

APPLICATION OF PATENT OF INVENTION


NUMBER	PRIORITY PARTICULARS DATE	COUNTRY
APPLICANT(S) QUÍMICA SINTETICA, S.A. ADDRESS: C. de l'Acer, 24 NATIONALITY: SPANISH 08004 BARCELONA, BARCELONA, SPAIN		
INVENTOR(S) ANTONIO COSME GOMEZ, JAVIER VILLASANTE PRIETO, FRANCISCO PALOMO NICOLAU		
INT. CL	GRAPHIC	
TITLE OF THE INVENTION		
CRYSTALLINE GATIFLOXACIN	FORM OF	
ABSTRACT CRYSTALLINE FORM OF GATIFLOXACIN <p>The present invention relates to a crystalline form of gatifloxacin obtainable by a process that comprises recrystallisation of the crude gatifloxacin in methanol and which is stable with a water content ranging between 2.5 and 4.5% by weight, to a process for preparing it and to the use thereof as an active substance in the preparation of pharmaceutical formulations.</p>		

CRYSTALLINE FORM OF GATIFLOXACIN**Field of the technique**

The present invention relates to a new
5 crystalline form of the active pharmaceutical substance
gatifloxacin.

Prior state of the art

Gatifloxacin is the international common name of
10 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-,
piperazinyl)-4-oxo-3-quinolinecarboxylic acid of formula
(I), with application in medicine and known for its
antibiotic activity:



European patent application EP-A-230295 discloses
20 the preparation of gatifloxacin, which is isolated in
hemihydrate form ($1/2 \text{ H}_2\text{O}$), corresponding to 2.34% in
calculated weight of water.

European patent application EP-A-805156 discloses
a sesquihydrated crystalline form ($3/2 \text{ H}_2\text{O}$), corresponding
25 to 6.72% in calculated weight of water.

Both crystalline forms have a tendency to absorb
water and to form polymorphs with a higher content in
hydration water.

Patent application WO-A-0222126 discloses
30 gatifloxacin pentahydrate ($5 \text{ H}_2\text{O}$), corresponding to 19.3%
in calculated weight of water.

There therefore exists a need to have another
hydrated crystalline form of gatifloxacin which is stable,
having a lower water content.

The authors of this invention have discovered a new crystalline form of gatifloxacin, which they have designated form I, which is stable with a water content ranging between 2.5 and 4.5% by weight when it is in
5 contact with the atmosphere at room temperature and with a relative humidity ranging between 20 and 70%.

Object of the invention

The object of the present invention is a new
10 crystalline form of gatifloxacin which is obtainable by means of a particular process.

Also object of this invention is the process for obtaining the new crystalline form of gatifloxacin.

Also forming part of the object of the present
15 invention is the use of the new crystalline form of gatifloxacin for the manufacture of a medicament for the treatment of infectious diseases of bacterial origin.

Brief description of the drawings

20 Figure 1 shows the powder X-ray diffractogram of the new crystalline form of gatifloxacin.

Figure 2 shows the powder X-ray diffractogram of the gatifloxacin hemihydrate taken from North American patent US5880283. Said patent includes only the X-ray
25 diffractogram, without the corresponding list of peaks shown at the different 2θ angles.

Figure 3 shows the ^{13}C nuclear magnetic resonance spectrum of the new crystalline form of gatifloxacin.

30 Detailed description of the invention

The authors of this invention have discovered a crystalline form of gatifloxacin, which they have designated form I, which is obtainable by means of a process comprising the following steps:

- the crude gatifloxacin is dissolved in methanol by heating to reflux temperature, using between 50 and 65 volumes of methanol for each unit by weight of crude gatifloxacin,
- 5 - it is cooled to a temperature between 15° C and 25° C within a period of time not exceeding 1.5 hours,
- during the cooling process it is seeded with form I gatifloxacin,
- it is then cooled to a temperature between 0° C and 5° C and kept at this temperature for at least 1 hour,
- 10 - the solid product obtained is separated by filtration, and
- the solid product is dried in an oven under vacuum to constant weight.

15 The crude gatifloxacin which is used as the starting product can be prepared as described in the Example of preparation set out below in this description, or according to the process described in Example 3 of the European patent application EP-A-230295.

20 The solution of crude gatifloxacin in methanol at reflux is prepared by using approximately 50 to 70 volumes of methanol for each unit by weight of crude gatifloxacin.

 Once the crude gatifloxacin has been dissolved at the reflux temperature of the methanol, the solution is cooled to a temperature ranging between 15° C and 25° C, which would be termed room temperature. In order to obtain the new crystalline form of gatifloxacin said cooling has to be carried out within a period of time not exceeding 1.5 hours.

30 During this period of cooling to room temperature seedings are carried out with form I gatifloxacin until a suspension containing an abundant precipitate is obtained.

 The form I gatifloxacin which is used for seeding for the first time is prepared by means of the process described in the Example of preparation set out below in

35

this description. On subsequent occasions the gatifloxacin obtained in Example 1 of this description can also be used.

The cooling of the suspension to a temperature between 0 and 5° C is carried out by means of refrigeration with cold water and it is kept at this temperature for approximately one hour.

The solid obtained is separated by filtration and washed with cold methanol.

The moist solid is dried in an oven at approximately 40° C *in vacuo* to constant weight.

The new crystalline form of gatifloxacin which is obtainable by this process has an initial water content ranging between 0.8 and 1.6% and stabilises with a water content ranging between 2.5 and 4.5% by weight, when it is in contact with the atmosphere at room temperature and with a relative humidity comprised between 20 and 70%, and it remains stable for at least 2 months in such conditions.

The new crystalline form of gatifloxacin usually stabilises in a period of time approximately equal to three days, but it could take longer to reach that degree of hydration if the degree of relative humidity was lower than 20%.

The crystalline form of gatifloxacin object of the present invention is characterised by its powder X-ray diffractogram (Figure 1), ¹³C nuclear magnetic resonance spectrum (Figure 3) and analysis of water content by the Karl-Fischer method.

The X-ray diffractogram of the hemihydrate (Figure 2) has been obtained from North American patent US5880283, in which the diffractograms of gatifloxacin hemihydrate (comparative substance) are compared with those of gatifloxacin sesquihydrate.

The form I gatifloxacin has an X-ray diffractogram which shows peaks at the 2θ angles 16.5 ± 0.2

and 17.8 ± 0.2 which are not present in the X-ray diffractogram of the gatifloxacin hemihydrate.

In its turn, the gatifloxacin hemihydrate has an X-ray diffractogram showing peaks at the 2θ angles 13.9 ± 0.2 , 14.5 ± 0.2 , 20.3 ± 0.2 , 22.5 ± 0.2 and 24.2 ± 0.2 , which are not present in the X-ray diffractogram of the form I gatifloxacin object of the invention.

For recording the powder X-ray diffractograms a PHILIPS X'Pert automatic diffractometer equipped with a Cu tube and a graphite secondary monochromator of the following technical specifications was used:

- Cu wavelength $K\alpha$: 1.5419 \AA ;
- receiving slit: 0.1 mm ;
- Soller: 0.04 radians ;
- 15 - dispersion slit and divergence slit: 1°

The tube worked at 40 kV and 50 mA . Sweeping was carried out continuously in the 2θ interval between 5 and 40° with pass of 0.03° and 1-second pass time.

The ^{13}C nuclear magnetic resonance spectrum (Figure 3) was recorded on a solid sample of form I gatifloxacin.

Also object of the invention is the process for preparing the new crystalline form of gatifloxacin, which comprises the following steps:

- 25 - the crude gatifloxacin is dissolved in methanol by means of heating to the reflux temperature,
- it is cooled to a temperature ranging between 15° C and 25° C in a period of time not exceeding 1.5 hours, and
- 30 - it is seeded with form I gatifloxacin.

The solution of crude gatifloxacin in methanol at reflux is preferably prepared by using between 50 and 70 volumes of methanol for each unit by weight of crude gatifloxacin.

Once the crude gatifloxacin has been dissolved at the reflux temperature of methanol, the solution is cooled to room temperature, preferably to a temperature ranging between 15 and 25° C. To prepare the new crystalline form of gatifloxacin this cooling has to be carried out within a period of time not exceeding 1.5 hours.

During this period of cooling to room temperature, seedings with form I gatifloxacin are carried out until a suspension containing an abundant precipitate is obtained.

The form I gatifloxacin which is used for the seeding the first time is prepared by means of the process described in the Example of preparation set out below in this description. On subsequent occasions the form I gatifloxacin obtained in Example 1 of this description can also be used.

The process for preparing the new crystalline form of gatifloxacin further comprises the following steps:

- the suspension is cooled to a temperature ranging between 0° C and 5° C and is kept at this temperature for at least 1 hour,
- the solid product is filtered, and
- the product is dried in an oven to constant weight.

The cooling of the suspension to a temperature between 0° C and 5° C is carried out by means of refrigeration with cold water and it is kept at this temperature for at least 1 hour.

The solid obtained is separated by filtration and washed with cold methanol.

The moist solid is dried in an oven, preferably at 40° C *in vacuo*, to constant weight.

The new crystalline form of gatifloxacin, which is obtainable by this process, has an initial water content ranging between 0.8 and el 1.6%, and as being in contact with the atmosphere at room temperature and with a relative

humidity ranging between 20 and 70%, it is stable with a water content between 2.5 and 4.5% by weight.

Surprisingly, it has been found that the new crystalline form of gatifloxacin containing between 2.5 and 4.5% of water by weight remains stable in its water content for at least 2 months, even at room temperature and with a relative humidity between 20 and 70%, and has excellent properties of disintegration and dissolution rate, which makes it very suitable for use as an active substance in pharmaceutical formulations, preferably for the manufacture of a medicament for the treatment of infectious diseases of bacterial origin.

The example which follows below is set out for the purposes of providing to the skilled man in the art a detailed explanation of a specific embodiment of the process for preparing the compound of the invention.

Example of preparation. Preparing form I gatifloxacin for seeding

10 10 g (0.0339 moles, 1 equivalent) of 1-cyclopropyl-6,7-difluoro-1,4-dihydro-8-methoxy-4-oxo-3-quinolinecarboxylic acid (CAS no.: 112811-72-0) is placed in a flask, 30 mL of acetonitrile (3 volumes) is added and the solution is heated to a temperature of 76-80° C. Once reflux has been attained, 3.28 g (0.0203 moles, 0.6 equivalents) of hexamethyldisilazane (HMDS) is added using a compensated addition funnel, maintaining the temperature. Once the addition is completed, the reaction is maintained with stirring for 1 hour at a temperature of 76-80° C.

30 Once this period has elapsed, the reaction mixture is cooled to a temperature between 0 and 15° C, and 5.78 g (0.0407 moles, 1.2 equivalents) of boron trifluoride ethyletherate is added, keeping the temperature below 15° C. Once the addition has finished, the temperature is

allowed to rise to 15-25° C and it is kept under these conditions for approximately 2 hours.

The pH of the mixture is then adjusted to an approximate value of 9 with triethylamine (approximately 2 mL). To the resulting suspension is added a solution of 10.19 g (0.1017 moles, 3 equivalents) of 2-methylpiperazine in 28 mL of acetonitrile, keeping the temperature between 15 and 25° C. The resulting amber solution is kept with stirring under these conditions for approximately 3 hours.

Once the reaction has been completed, the mixture is distilled at low pressure until a stirrable paste is obtained. At this point, 50 mL of methanol is added, the resulting suspension is raised to a temperature of 63-67° C and kept under these conditions for approximately 5 hours. Once the reaction has been completed the mixture is cooled to a temperature of 25-35° C over a water bath and then to a temperature of 0-5° C over a water/ice bath for a further 1 hour. The resulting precipitate is filtered, washed with cold methanol (2 x 10 mL) and dried at 40° C in an oven in vacuo to constant weight. 10.70 g of crude gatifloxacin is obtained, with a water content of 2.95% by weight. The yield of the process is 81.8%.

The crude product obtained is used for the seeding in Example 1.

The crude product is crystallised in methanol by dissolving 20 g of crude gatifloxacin in 1 l of methanol (50 volumes) at a temperature of 63-67° C. Once all the product has been dissolved it is placed to cool to a temperature of 30-40° C, and then to a temperature of 0-5° C over a water/ice bath, maintaining it under these conditions for 1 hour. The resulting suspension is filtered and the solid retained is washed with 20 mL (1 volume) of cold methanol. The solid obtained is dried at 40° C in a vacuum oven to obtain 18.65 g of gatifloxacin with a water content of 2.36% by weight.

The product obtained is used as crude gatifloxacin (starting product).

Example 1.- Preparing form I gatifloxacin

5 To 39.23 g of crude gatifloxacin, prepared as in
the Example of preparation, are added 2 l of methyl
alcohol, and the suspension is heated to reflux
temperature. Once a reflux regime has been reached, methyl
alcohol is added until the product has totally dissolved.
10 The total volume of methyl alcohol used is 2.69 l. Once the
product has been dissolved the solution is cooled to room
temperature in one hour, carrying out seedings with form I
gatifloxacin (obtained in the Example of preparation) until
the amount used in the seeding is inconsiderable compared
15 with the amount of crystallised product. The resulting
suspension is then cooled to a temperature between 0 and 5°
C over a water/ice bath, and is kept within that
temperature range for 1 hour. Once the cooling period has
elapsed, the solid is isolated by filtration and washed
20 with cold methyl alcohol (2 x 40 mL). The product obtained
is dried in a vacuum oven at 40° C to constant weight.

31.25 g of a white solid is obtained, which has a
water content, at the time of taking it out of the
desiccator, of 1.5% by weight. The yield is 80.8%.

25 The product thus obtained is kept at room
temperature in contact with the atmosphere, and three days
later it has a water content of 3.22% by weight, which
remains stable for at least 2 months at room temperature
and with a relative humidity between 20 and 70%.

30 Table 1 shows the water-content values of the
form I gatifloxacin obtained during the stability test:

Table 1

Time	0	3 hours	6 hours	3 days	13 days	17 days	68 days
% water by weight	1.5	2.2	2.3	3.2	2.9	3.0	3.3

The powder X-ray diffractogram recorded on the
5 sample of form I gatifloxacin remains substantially
unchanged over this entire period of time.

CLAIMS

1. A crystalline form of gatifloxacin obtainable by means of a process which includes the following steps:

- 5 - the crude gatifloxacin is dissolved in methanol by heating to reflux temperature, using between 50 and 65 volumes of methanol for each unit by weight of crude gatifloxacin,
- 10 - it is cooled to a temperature ranging between 15° C and 25° C within a period of time not exceeding 1.5 hours,
- during the cooling process it is seeded with form I gatifloxacin,
- 15 - it is then cooled to a temperature comprised between 0° C and 5° C and kept at this temperature for at least 1 hour,
- the solid product is separated by filtration, and
- the solid product is dried in an oven under vacuum to constant weight.

20 2. A process for preparing a crystalline form of gatifloxacin, which includes the following steps:

- the crude gatifloxacin is dissolved in methanol by means of heating to the reflux temperature,
- 25 - it is cooled to a temperature between 15° C and 25° C in a period of time not exceeding 1.5 hours, and
- it is seeded with form I gatifloxacin.

30 3. A process according to Claim 2, characterised in that the crude gatifloxacin is dissolved in methanol, using between 50 and 70 volumes of methanol for each unit by weight of crude gatifloxacin.

4. A process according to Claims 2 and 3, characterised in that it further comprises the following steps:

- the suspension is cooled to a temperature between 0° C and 5° C and kept at that temperature for at least 1 hour,
- the solid product is filtered, and
- 5 - the product is dried in an oven to constant weight.

5. Use of the form of gatifloxacin according to Claim 1 for the manufacture of a medicament for the treatment of infectious diseases of bacterial origin.

ABSTRACT

The present invention relates to a crystalline form of gatifloxacin obtainable by a process that comprises
5 recrystallisation of the crude gatifloxacin in methanol and which is stable with a water content ranging between 2.5 and 4.5% by weight, to a process for preparing it and to the use thereof as an active substance in the preparation of pharmaceutical formulations.